arXiv:1205.4354v1 [math.FA] 19 May 2012

DIRECTLY FINITE ALGEBRAS OF PSEUDOFUNCTIONS ON LOCALLY COMPACT GROUPS

YEMON CHOI

March 4, 2013

Abstract

An algebra A is said to be directly finite if each left invertible element in the (conditional) unitization of A is right invertible. It has long been known that the complex group algebra of a discrete group is directly finite. We extend this result, using some Hilbert algebra techniques, and show that the reduced group C*-algebra of a unimodular group is directly finite.

We then investigate the corresponding problem for algebras of p-pseudofunctions, where $p \neq 2$. We show that these algebras are directly finite if G is amenable and unimodular, or unimodular with the Kunze-Stein property. In contrast, we show that $L^1(G)$ is not directly finite if G is the affine group of either the real or complex line.

MSC 2010: 22D15, 43A15 (primary); 22D25, 46L05 (secondary)

1 INTRODUCTION

MOTIVATION AND OVERVIEW

Direct finiteness of an algebra has been defined in the abstract. Our original motivation for investigating the direct finiteness of group algebras comes from the following general question. Let G be a locally compact group, let $1 < p < \infty$, and let $f \in C_c(G)$; then, by integrating f against the left regular representation of G on $L^p(G)$, we obtain a bounded linear operator $\lambda_p(f) : L^p(G) \to L^p(G)$. *How big is the approximate point spectrum of* $\lambda_p(f)$?

If G is abelian, then by using Gelfand/Fourier theory we find that the spectrum of $\lambda_p(f)$ consists entirely of approximate eigenvalues; that is, the approximate point spectrum is as big as possible. In the non-abelian context new tools are needed. It seems to be part of the folklore (see, for instance, the author's article [3]) that when G is discrete, the spectrum of $\lambda_2(f)$ consists entirely of approximate eigenvalues, for all $G \in c_{00}(G)$. The key result needed for the proof is an old observation, apparently first noted by Kaplansky, that the group algebra of a *discrete* group is directly finite. (See [3, Remark 2.10] for some further remarks on the history of this observation, and later proofs.)

Both Kaplansky's proof and the later one of Montgomery [15] yield more: they shows that the reduced group C*-algebra of a discrete group is directly finite. It is therefore natural to ask if the same is true for an arbitrary locally compact group. One of our main results is that the reduced group C*-algebra of any *unimodular* group is directly finite (Theorem 3.6 below). It seems likely that this result can be used to obtain results analogous to those in the author's previous article [3]; we intend to treat this more fully in forthcoming work, and so will not discuss these implications here.

It is natural to ask if the same is true for $\mathrm{PF}_p(G)$, the algebra of p-pseudofunctions on G, when $p \neq 2$. We have not been able to obtain a proof or a counter-example for general unimodular groups, but by building on the C^*-case we obtain partial results. Namely, $\mathrm{PF}_p(G)$ is directly finite if G is unimodular and also has one of the following two properties: amenability, or the so-called "Kunze-Stein" property. Examples of the latter case are semisimple Lie groups with finite centre, and certain automorphism groups of trees. (We note that a group is both amenable and Kunze-Stein if and only if it is compact.) The cases where G is amenable and unimodular follow easily from an embedding result of Herz: since the result is key to our work, and since the reader who wishes to learn the proof from Herz's papers must face many technical details related to but not needed for the embedding theorem, we have included further details in an appendix.

It turns out that we cannot drop the assumption of unimodularity. Somewhat to the author's surprise, there are solvable Lie groups for which $L^1(G)$, and hence $\mathrm{PF}_p(G)$ for every p, are not directly finite. The precise examples are given below in Theorem 5.1. The proof combines a theorem of Leptin, which says that the inclusion $L^1(G) \to \mathrm{C}^*_r(G)$ is spectrum-preserving when G is either of the groups described above, with calculations of Diep and of Rosenberg that were used in *classifying* $\mathrm{C}^*_r(G)$ for certain solvable Lie groups G.

REMARKS ON THE APPROACH OF THE PRESENT PAPER

Some of these results presented here were included in an old preprint of the author, "*Group* C^**-algebras which are quasi-directly finite*", arXiv `1003.1650`. That preprint, which remains unpublished, is now superseded by the present article.

We have attempted to make this paper accessible to those working in areas such as operator theory, Banach algebras, or non-abelian harmonic analysis, who like the author are not specialists in the theory of operator algebras. This has meant including material that will be well known to such specialists, either to provide extra background, or to provide references to sources where the reader can find actual proofs (as opposed to assertions or references to proofs). Our aim was not to shun citations when citation would suffice. Rather, we wished to avoid referring to sources where a reader might have to digest large amounts of unnecessary technical material before getting to the particular parts needed to follow the present paper.

Moreover, in the interests of accessibility, we sometimes avoid giving the most general definitions and results; in particular, we adopt a relatively "low-tech" approach to the definition of $\mathrm{PF}_p(G)$, rather than using the full machinery that would be needed for serious study of general convolution operators on $L^p(G)$. Likewise, when we need to appeal to the theory of Hilbert algebras, we shall avoid any discussion of left Hilbert algebras.

BACKGROUND ON CONVOLUTION OPERATORS AND PSEUDOFUNCTIONS

We close this introductory section with a short account of what is needed to define and work with $\mathrm{PF}_p(G)$. This material is included because in other sources, the initial preliminaries are either summarized without explanation, or done at a more technical level than is needed for our purposes. It seems worthwhile to give the reader a less comprehensive and more simple-minded approach, which nevertheless suffices for the present paper.

For additional details and further references, a recent source is the monograph [5], although the reader should note that some of the basic properties are stated without full proofs. The paper [10] is also invaluable reading for those who wish to study convolution operators on $L^p(G)$ in greater depth.

Let us fix some notation and terminology. Let G be a locally compact group, equipped with a fixed left Haar measure μ. We denote by $C_c(G)$ the space of continuous, compactly supported, complex-valued functions on G; this is an algebra when equipped with convolution as product. For each $p \in (1,\infty)$, define $\lambda_p : C_c(G) \to \mathcal{B}(L^p(G))$ by

$$\lambda_p(f)(h) = f * h \qquad f \in C_c(G), h \in L^p(G).$$

The norm-closure of $\lambda_p(C_c(G))$ inside $\mathcal{B}(L^p(G))$ is a closed subalgebra, which we denote by $\mathrm{PF}_p(G)$ and call the algebra of p-pseudofunctions on G. Note that $\mathrm{PF}_2(G)$ is nothing but the reduced group C^*-algebra of G.

REMARK 1.1. Using standard results about integration on locally compact groups, one can show λ_p extends to an *injective*, continuous algebra homomorphism $L^1(G) \to \mathrm{PF}_p(G)$. However, we only need to mention this at one point (Remark 5.2), and it plays no significant role in our main arguments.

2 DIRECTLY FINITE BANACH ALGEBRAS

We start with some basic results that will be needed later, but which may also be of some independent interest. Throughout, we adopt the convention that a ring need not be commutative, nor contain a multiplicative identity.

A ring R with identity is said to be directly finite if each left-invertible element of R is invertible. Motivated by examples from semigroup theory, Munn [16] generalized this definition to the setting of rings without identity. It is convenient to present this definition and some basic consequences by using the notions of left and right quasi-inverses. (In the Banach-algebraic setting, a convenient reference is e.g. [2, §4].)

DEFINITION 2.1. Let R be a ring. Given $a, b \in R$, let $a \bullet b := a + b - ab$. If $a \bullet b = 0$ then we say that a is a left quasi-inverse for b and b is a right quasi-inverse for a. An element which has both a left and a right quasi-inverse is said to be quasi-invertible.

The definition of $\bullet$ is cooked up to ensure that if R has an identity element 1, then

$$1 - a \bullet b = (1-a)(1-b). \tag{1}$$

and it is clear from this, or directly from the definition, that $a \bullet 0 = a = 0 \bullet a$ for all $a \in A$. Moreover, the operation $\bullet$ is associative: one could check this by a direct calculation, but it is more instructive to adjoin a formal identity 1 and observe, using (1) repeatedly, that

$$\begin{aligned} 1 - (a \bullet b) \bullet c = (1 - a \bullet b)(1-c) &= (1-a)(1-b)(1-c) \\ &= (1-a)(1 - b \bullet c) = 1 - a \bullet (b \bullet c) \end{aligned}$$

REMARK 2.2. It is more intuitive to reason with left, right and two-sided invertible elements than with their "quasi-" counterparts. On the other hand, the language of quasi-inverses streamlines the statements of some results, and in particular avoids clunky use of conditional unitizations.

DEFINITION 2.3. A ring R is directly finite if any pair $(a,b) \in R \times R$ satisfying $a \bullet b = 0$ also satisfies $b \bullet a = 0$.

REMARK 2.4. In [16], a ring with this property is said to be "quasidirectly finite". We have chosen to extend the terminology from the unital case to the non-unital one, rather than introduce slightly cumbersome extra terminology.

The property of being directly finite is clearly inherited by arbitrary subrings (even if our larger ring has an identity, we do not require that subrings contain this identity). Consequently, if S is a directly finite ring and $\phi : R \to S$ is an injective homomorphism, R is directly finite; this holds even if R and S have identity elements and $\phi(1_A) \neq 1_S$.

At this point, since we will need to briefly discuss unitizations, we switch to algebras rather than rings. (This is solely to sidestep the annoyance that the ring-unitization of an algebra is not its algebra-unitization; one could formulate and prove the analogous results for ring-unitizations just as easily.) Recall that the forced unitization of a k-algebra A is the vector space $A \oplus k1$, where 1 is a formal identity element, equipped with multiplication in the natural way. We denote the forced unitization of A by $A^\sharp$.

LEMMA 2.5.

(i) *Let A be an algebra (with or without identity). Then A is directly finite (in the sense of Definition 2.3) if and only if each left-invertible element of $A^\sharp$ is invertible.*

(ii) *Let A be a ring with identity. Then A is directly finite (in the sense of Definition 2.3) if and only if each left-invertible element of A is invertible.*

The lemma follows easily from the identity (1); we omit the proof.

In a unital Banach algebra, the group of invertible elements is open in the norm topology. There is an analogous result for quasi-inverses, see [2, Theorem 4.8]; we shall require a slightly more precise version.

LEMMA 2.6. *Let A be a Banach algebra and let $c \in A$. Suppose there exists $b \in A$ with $b \bullet c = 0$. Then, for each $c' \in A$ that is sufficiently close to c, there exists $a \in A$ such that $a \bullet b \bullet c' = 0$.*

Proof. Give the unitization $A^\sharp$ its usual norm, so that $\|1\| = 1$. We have $(1-b)(1-c) = 1$ in $A^\sharp$. Let $\delta = (1 + \|b\|)^{-1} > 0$; then, given any $c' \in A$ such that $\|c' - c\| < \delta$, define

$$u := (1-b)(1-c') \in A^\sharp.$$

Since $1 - u = (1-b)(c'-c)$ has norm < 1, u is invertible in the Banach algebra $A^\sharp$, and

$$a := 1 - u^{-1} = -\sum_{n\geq 1} (1-u)^n$$

lies in A. By construction, $1 - u = b \bullet c'$, so that $a \bullet b \bullet c = 0$ as required. ☐

The following proposition is the main technical result of this section. At one point later, we will need the extra generality afforded by left ideals, rather than two-sided ones.

PROPOSITION 2.7. *Let A be a Banach algebra, and let J be a* left *ideal in A which is dense for the norm topology. Then J is directly finite if and only if A is.*

Proof. Clearly, if A is directly finite then so is J. Conversely, suppose J is directly finite, and let $b, c \in A$ satisfy $b \bullet c = 0$. By Lemma 2.6 and density of J in A, we can find $c' \in J$ and $a \in A$ such that $a \bullet b \bullet c' = 0$.

Put $b' = a \bullet b$; then since $b' \bullet c' = 0$, we have $b' = b'c' - c' \in J$, as $c' \in J$ and J is a left ideal. Since J is directly finite, $c' \bullet b' = 0$, that is, $c' \bullet a \bullet b = 0$. So b has a left quasi-inverse in A, and as $b \bullet c = 0$ we conclude that $c \bullet b = 0$ as required. ☐

3 DIRECTLY FINITE C^*-ALGEBRAS, VIA DENSELY-DEFINED TRACES

We start by fixing some notation and terminology, which we take from [19, §5]. If A is a C^*-algebra, A_+ will denote its cone of positive elements. A weight on A_+ is a function $\phi : A^+ \to [0, \infty]$ that is $\mathbb{R}_+$-linear and additive. ϕ is faithful if $\phi(x) > 0$ for all $x \in A_+ \setminus \{0\}$; it is tracial if it satisfies $\phi(u^*au) = \tau(u)$ for all $a \in A_+$ and all unitary $u \in A^\sharp$.

Given a tracial weight τ on A_+, there exists a 2-sided $*$-ideal $A^\tau \subseteq A$, and a linear tracial functional $A^\tau \to \mathbb{C}$ which coincides with τ on $A^\tau \cap A^+$; by abuse of notation, we will denote this functional also by τ. A^τ is called the ideal of definition of the trace τ. Moreover, the set $A_2^\tau := \{x \in A : \tau(xx^*) < \infty\}$ has the same norm-closure in A as does A^τ. (See [19, Propositions 5.1.2 and 5.2.2] and their proofs for the details.)

PROPOSITION 3.1. *Let B be a unital* C^**-algebra and τ a faithful tracial weight on B_+. Let A denote the norm-closure of B^τ inside B. Then A is directly finite.*

When B is unital and $B^\tau = B$, this result is known, and is used in [15] to prove that the complex group algebra of a (discrete) group is always directly finite. The author is unaware of a reference which explicitly states Proposition 3.1 in the generality given here, so a full proof will be given below. The key observation is the following standard result about C^*-algebras, which seems to be part of the folklore.

LEMMA 3.2. *If p is an idempotent in a* C^**-algebra, there exists a hermitian idempotent e in that algebra which satisfies $ep = p$ and $pe = e$.*

The usual, somewhat post-hoc proof of the lemma is as follows: after adjoining an identity element to our algebra if necessary, we make the *Ansatz*

$$e = pp^*(1 + (p - p^*)(p^* - p))^{-1}, \qquad (\dagger)$$

which clearly satisfies $e = pe$. It remains to check that $ep = p$ and that e is an idempotent. Here is one way: take a faithful representation of our (unital) C^*-algebra on a Hilbert space H, and regard p as a 2×2 operator matrix $\begin{pmatrix} I & R \\ 0 & 0 \end{pmatrix}$ with respect to the decomposition $H = \operatorname{ran}(p) \oplus \operatorname{ran}(p)^\perp$. Computing the right-hand side of (†) with respect to this block-matrix decomposition gives $\begin{pmatrix} I & 0 \\ 0 & 0 \end{pmatrix}$, which is the orthogonal projection of H onto $\operatorname{ran}(p)$. An alternative approach, taken in [11], is to exploit the theory of polynomial identities; we thank J. Meyer [14] for bringing this to our attention.

Proof of Proposition 3.1. By Proposition 2.7, it suffices to show that B^τ is directly finite. Let $a, b \in B^\tau$ be such that $a \bullet b = 0$, and let $p := b \bullet a = ab - ba \in B^\tau$. If we can show $p = 0$ then we will be done.

Observe that p is an idempotent, since

$$2p - p^2 = p \bullet p = b \bullet a \bullet b \bullet a = b \bullet 0 \bullet a = p\,.$$

By Lemma 3.2, there is a self-adjoint idempotent $e \in A$ such that $pe = e$ and $ep = p$; in particular, $e \in B^\tau$, since B^τ is a (right) ideal in A. The tracial property of τ then implies

$$\tau(e^*e) = \tau(e) = \tau(pe) = \tau(ep) = \tau(p) = \tau(ab) - \tau(ba) = 0\,.$$

As τ is faithful, this forces $e = 0$, and hence forces $p = 0$, as required. ☐

REMARK 3.3. When using Proposition 3.1 to prove that a given C*-algebra A is directly finite, it is sometimes easier or more convenient to create the tracial weight τ on a larger containing algebra B, and then to show that $B^\tau \cap A$ is dense in A.

A particular case of Proposition 3.1 arises when B is taken to be the left (or right) von Neumann algebra associated to a Hilbert algebra. Recall that a Hilbert algebra is an associative, complex $*$-algebra $\mathfrak{A}$, equipped with an inner product $(\cdot \mid \cdot)$ that satisfies certain compatibility conditions: see [7, ch. I§5, défn 1] or [18, Definition 11.7.1]. Completing $\mathfrak{A}$ in the norm $x \mapsto (x \mid x)^{1/2}$ yields a Hilbert space $\mathcal{H}$, equipped with natural left and right actions of $\mathfrak{A}$, called the Hilbert space associated to $\mathfrak{A}$. Given $a \in \mathfrak{A}$ we denote by $U_a : \mathcal{H} \to \mathcal{H}$ the unique operator satisfying $U_a(x) = ax$ for all $x \in \mathfrak{A}$; the map $a \mapsto U_a$ is an injective $*$-homomorphism from $\mathfrak{A}$ into $\mathcal{B}(\mathcal{H})$. The weak-operator closure of $\{U_a \; : \; a \in \mathfrak{A}\}$ is denoted by $\mathcal{U}(\mathfrak{A})$, and called the left von Neumann algebra associated to $\mathfrak{A}$. (A summary of these and related results can be found in [18, §11.7]; full details and proofs are given in [7, ch. I,§§5–6].)

THEOREM 3.4. *Let $\mathfrak{A}$ be a Hilbert algebra, and let $\mathfrak{B}$ be the norm-closed subalgebra of $\mathcal{U}(\mathfrak{A})$ generated by the set $\{U_a \; : \; a \in \mathfrak{A}\}$. Then $\mathfrak{B}$ is directly finite.*

Proof. We use freely the concepts discussed in [7, ch. I, §§5–6], to which the reader is referred for unexplained terminology.

Let $\mathfrak{n}$ be the set of all $T \in \mathcal{U}(\mathfrak{A})$ that are of the form $T = U_a$ for some $a \in \mathcal{H}$ that is bounded relative to $\mathfrak{A}$ [7, ch. I, §5, défns 2 and 3]. Then $\mathfrak{n}$ is a 2-sided ideal in $\mathcal{U}(\mathfrak{A})$ [7, ch. I, §5, prop. 3], and $\{U_a \; : \; a \in \mathfrak{A}\} \subseteq \mathfrak{n}$. Define $\varphi : \mathcal{U}(\mathfrak{A})^+ \to [0,\infty]$ by

$$\varphi(S) = \begin{cases} (a \mid a) & \text{if } S^{1/2} = U_a \in \mathfrak{n}\,, \\ +\infty & \text{otherwise.} \end{cases}$$

One can show φ is a faithful, tracial weight on $\mathcal{U}(\mathfrak{A})$ (the so-called canonical trace associated to $\mathfrak{A}$).

Let $\mathfrak{m}_\varphi \subseteq \mathcal{U}(\mathfrak{A})$ denote the ideal of definition of φ. It suffices to show that $\mathfrak{m}_\varphi \cap \mathfrak{B}$ is norm-dense in $\mathfrak{B}$; therefore, since each element of $\mathfrak{B}$ is a linear combination of (at most four) elements in its positive cone, it suffices to show that $\mathfrak{B}_+ \cap \mathfrak{m}_\varphi$ is dense in $\mathfrak{B}_+$. Given $b \in \mathfrak{B}^+$, we know $b^{1/2} \in \mathfrak{B}^+$, so there is a sequence $(a(n))_{n\geq 1} \subset \mathfrak{A}$ satisfying $U_{a(n)} \to b^{1/2}$ in norm, and hence satisfying $U^*_{a(n)} U_{a(n)} \to b$ in norm. As $U^*_a = U_{(a^*)} \in \mathfrak{n}$ for all $a \in \mathfrak{A}$, $U_{a^*a} = U^*_a U_a \in \mathfrak{B}_+ \cap \mathfrak{m}_\varphi$ for all $a \in \mathfrak{A}$, so we have an approximating sequence of the desired form. ☐

REMARK 3.5. It is important in Theorem 3.4 that we take the norm-closure of the Hilbert algebra $\mathfrak{A}$, and not its strong operator closure. For instance, take $\mathfrak{A}$ to be the algebra of finite rank operators on ℓ^2, with $(S \mid T)$ defined to be the trace of TS. Then $\mathfrak{H}$ can be identified with ℓ^2, and $\mathfrak{B}$ is the algebra $\mathcal{K}(\ell^2)$ of compact operators, which is directly finite (this follows of course from our theorem, but more generally $\mathcal{K}(X)$ is directly finite for any Banach space X, as observed in [3, §3]). On the other hand, $\mathfrak{A}$ is dense in $\mathcal{B}(\ell^2)$ for the strong operator topology, and $\mathcal{B}(\ell^2)$ is clearly not directly finite.

We now apply Theorem 3.4 to certain group C^*-algebras.

THEOREM 3.6. *If G is unimodular, then $\mathrm{C}^*_r(G)$ is directly finite.*

Proof. Let $\mathfrak{A}$ denote the space $C_c(G)$ of continuous, compactly supported functions on G. We equip $\mathfrak{A}$ with its usual involution $f^*(x) := \overline{f(x^{-1})}$ $(f \in \mathfrak{A}, x \in G)$ and the inner product $(f \mid g) = \int_G f(x)\overline{g(x)}\, d\mu(g)$ $(f, g \in \mathfrak{A})$. It is straightforward to check that $\mathfrak{A}$ is a Hilbert algebra, see [7, chapitre 1, §5, exercice 5] and the ensuing hints, or [8, §13.10], or [18, Example 11.7.2]. (This relies crucially on G being unimodular.) The associated Hilbert space $\mathcal{H}(\mathfrak{A})$ is just $L^2(G)$; the associated (left) von Neumann algebra $\mathcal{U}(\mathfrak{A})$ is $\mathrm{VN}(G)$, the usual group von Neumann algebra of G; and the norm-closure of $\mathfrak{A}$ inside $\mathcal{U}(\mathfrak{A})$ is $\mathrm{C}^*_r(G)$. Applying Theorem 3.4, the result follows. ☐

It is natural to ask if the converse of Theorem 3.6 holds. The following examples show it does not, although in some sense they constitute a 'cheat answer'.

EXAMPLE 3.7 (A certain family of solvable Lie groups). Let p, q be strictly positive integers, and let $\alpha = (\alpha_1, \dots, \alpha_{p+q})$ where each α_j is a strictly positive real number. Denote by $G(p, q, \alpha)$ the semidirect product $\mathbb{R}^{p+q} \rtimes \mathbb{R}$, where the action of $\mathbb{R}$ on $\mathbb{R}^{p+q}$ is given by

$$t \mapsto \mathrm{diag}(e^{\alpha_1 t}, \dots, e^{\alpha_p t}, e^{-\alpha_{p+1} t}, \dots, e^{-\alpha_{p+q} t})$$

In general, $G(p, q, \alpha)$ is not unimodular. However, it turns out that the isomorphism class of $\mathrm{C}^*(G(p, q, \alpha))$ depends only on p and q and not on α. (See [21, pp. 12–13] and [20, pp. 190–191].) Moreover, if we happen to choose α such that $\alpha_1 + \cdots + \alpha_p = \alpha_{p+1} + \cdots + \alpha_{p+q}$, then $G(p, q, \alpha)$ *is* unimodular; see [20, p. 190]. Therefore, by Theorem 3.6, $\mathrm{C}^*_r(G(p, q, \alpha))$ is directly finite.

4 SOME GROUPS FOR WHICH $\mathrm{PF}_p(G)$ IS DIRECTLY FINITE

Recall that the reduced group C^*-algebra of G is the case $p = 2$ of the algebra $\mathrm{PF}_p(G)$

One would like to generalize Theorem 3.6 to cover $\mathrm{PF}_p(G)$ for G unimodular and all $p \in (1,\infty)$, or to find a unimodular example for which $\mathrm{PF}_p(G)$ is not directly finite for some $p \neq 2$. While we have been unable to do this, there are partial results we can obtain that apply to two different classes of groups.

REMARK 4.1. Let G be a unimodular, locally compact group. Let p and q be conjugate indices, strictly between 1 and ∞. The operator $\mathcal{B}(L^p(G)) \to \mathcal{B}(L^q(G))$, $T \mapsto T^*$, is a conjugate-linear anti-isomorphism, which maps $\mathrm{PF}_p(G)$ onto $\mathrm{PF}_q(G)$. It follows that $\mathrm{PF}_{p'}(G)$ is directly finite if and only if $\mathrm{PF}_p(G)$ is.

REMARK 4.2. Suppose we have two algebra norms on a given algebra A, say $\|\cdot\|_m$ and $\|\cdot\|_M$, such that $\|a\|_m \leq \|a\|_M$ for all $a \in A$. If A_m and A_M denote the corresponding completions of A, then of course there is a continuous algebra homomorphism $A_M \to A_m$, but this homomorphism need not be injective (just think of the maximal and the reduced C^*-algebra norms on $C_c(G)$.) Consequently, even if A_m is directly finite, extra work may be needed to show that A_M is.

We start by considering the case where G is amenable. The following theorem is a weaker version of one due to Herz.

THEOREM 4.3 (Herz). *Let G be a locally compact, amenable group. Regard λ_p as a homomorphism from $C_c(G)$ to $\mathrm{PF}_p(G)$. Then there is an injective homomorphism $J : \mathrm{PF}_p(G) \to \mathrm{PF}_2(G)$ such that $\lambda_2 = J\lambda_p$.*

Combining Theorem 4.3 with Theorem 3.6, the following is immediate.

COROLLARY 4.4. *Let G be an amenable, unimodular, locally compact group. Then $\mathrm{PF}_p(G)$ is directly finite, for every $1 < p < \infty$.*

REMARK 4.5. Herz actually proved something rather stronger. Let $\mathrm{CV}_p(G)$ denote the subalgebra of $\mathcal{B}(L^p(G))$ that consists of all operators which commute with right translations. It is straightforward to check that $\mathrm{CV}_p(G)$ contains the ultraweak closure of $\mathrm{PF}_p(G)$, which is itself an algebra, usually denoted by $\mathrm{PM}_p(G)$. It was shown in [9, 10] that when G is amenable, $\mathrm{PM}_p(G) = \mathrm{CV}_p(G)$ and we have a unital embedding $\mathrm{PM}_p(G) \to \mathrm{VN}(G)$ that extends $\lambda_2 : C_c(G) \to \mathrm{VN}(G)$.

Clearly, this embedding result implies Theorem 4.3: however, extracting a complete proof from these papers requires some work, and the task is made more arduous since both papers are dense and contain other highly technical results. Therefore, in Appendix A, we have included details on how to one can extract from [9, 10] the bare minimum needed to prove Theorem 4.3.

DEFINITION 4.6. A locally compact group G is said to have the Kunze-Stein property, or to be a Kunze-Stein group, if for each $1 \leq p < 2$ there exists a constant $C_p \geq 1$ such that

$$\|g * h\|_2 \leq C_p \|g\|_p \|h\|_2 \quad \text{for all } g \in L^p(G) \text{ and all } h \in C_c(G). \tag{2}$$

We mention two families of unimodular groups with the Kunze-Stein property.

(i) Let G be a connected semisimple Lie group with finite centre, such as $SL(n,\mathbb{R})$. Cowling [4] proved that G has the Kunze-Stein property. (The particular case $n = 2$ was established by Kunze and Stein in [12].)

(ii) Let T be a homogeneous tree of order ≥ 3, and equip $\mathrm{Aut}(T)$ with the topology of pointwise convergence; this makes it into a locally compact, totally disconnected group. Let G be a closed subgroup of $\mathrm{Aut}(T)$ which acts transitively on the boundary ∂T. (For instance, when $\mathbb{K}$ is a local field, the group $\mathrm{SL}(2, \mathbb{K})$ is isomorphic as a topological group to one of this form.) Then G has the Kunze-Stein property [17].

It is also remarked in [4] that the only amenable groups with the Kunze-Stein property are the compact ones. In view of this, the following result is a somewhat surprising counterpart to Corollary 4.4.

THEOREM 4.7. *Let G be unimodular and Kunze-Stein. Then $\mathrm{PF}_p(G)$ is directly finite.*

The proof of Theorem 4.7 occupies the rest of this section. The main idea is to define a dense, one-sided ideal in $\mathrm{PF}_p(G)$ which is also a subalgebra of $\mathrm{PF}_2(G)$, and then use Theorem 3.6. Morally speaking, for $1 \leq p < 2$ we wish to work with $\mathrm{PF}_p(G) \cap L^p(G)$; however, as we have defined things, this intersection is meaningless since $\mathrm{PF}_p(G)$ is a space of operators on $L^p(G)$ and not a space of functions or measures on G. We could make this precise and meaningful if we viewed both p-pseudofunctions and L^p functions as distributions on G, or as suitable limits of finite measures on G. Instead, we prefer to take an indirect approach, which is less concise but requires less machinery.

Define $\mathrm{diag}_p : C_c(G) \to L^p(G) \oplus_1 \mathrm{PF}_p(G)$ to be the 'diagonal' embedding

$$\mathrm{diag}_p(f) = (f, \lambda_p(f))$$

and define $X_p(G)$ to be the norm closure of $\mathrm{diag}_p(C_c(G))$. Let π_L and π_P denote the coordinate projections $L^p(G) \oplus_1 \mathrm{PF}_p(G) \to L^p(G)$ and $L^p(G) \oplus_1 \mathrm{PF}_p(G) \to \mathrm{PF}_p(G)$.

LEMMA 4.8. *The restrictions of π_L and π_P to the subspace $X_p(G)$ are both injective.*

Proof. If $(g, T) \in X_p(G)$ and $h \in C_c(G)$, an easy continuity argument shows that $g * h = T(h) \in L^p(G)$. Moreover, if $S \in \mathcal{B}(L^p(G))$ and $S(h) = 0$ for all $h \in C_c(G)$, then $S = 0$. The result follows. ☐

LEMMA 4.9.

(i) *Let $S \in \mathrm{PF}_p(G)$ and $(g, T) \in X_p(G)$. Then $(S(g), ST) \in X_p(G)$.*

(ii) *The bilinear map $X_p(G) \times X_p(G) \to X_p(G)$, defined by*

$$((f, S), (g, T)) \mapsto (S(g), ST)$$

is associative, and makes $X_p(G)$ a Banach algebra; moreover, $\pi_P : X_p(G) \to \mathrm{PF}_p(G)$ is an algebra homomorphism with dense range.

Proof. For this proof, we denote the usual L^p-norm on $C_c(G)$ by $\|\cdot\|_p$, anddenote the operator norm on $\mathcal{B}(L^p(G))$ by $\|\cdot\|_{p\to p}$.

Let (g_n) be a sequence in $C_c(G)$ with $\mathrm{diag}(g_n) \to (g, T)$; that is,

$$\|g_n - g\|_p \to 0 \quad \text{and} \quad \|\lambda_p(g_n) - T\|_{p\to p} \to 0$$

Let (f_n) be a sequence in $C_c(G)$ with $\|\lambda_p(f_n) - S\|_{p\to p} \to 0$. Then $f_n * g_n \in C_c(G)$; and a standard "3ε argument" shows that $\|f_n * g_n \to S(g)\|_p \to 0$ and $\|\lambda_p(f_n * g_n) - ST\|_{p\to p} \to 0$. This proves part (i).

For part (ii): associativity can be checked directly, as can the fact that the norm on $X_p(G)$ is submultiplicative. So $X_p(G)$ is a Banach algebra. Clearly $\pi_P : X_p(G) \to \mathrm{PF}_p(G)$ is a homomorphism; it has dense range, since $\pi_p \operatorname{diag}_p = \lambda_p$. ☐

LEMMA 4.10. *When G is Kunze-Stein, there exist injective, continuous linear maps* $\imath_{\mathrm{KS}} : L^p(G) \to \mathrm{PM}_2(G)$ *and* $\widetilde{\lambda}_{p,2} : X_p(G) \to \mathrm{PF}_2(G)$ *which make the following diagram commute:*

$$\begin{tikzcd}
 & & L^p(G) \arrow[dd, "\imath_{\mathrm{KS}}"] \\
C_c(G) \arrow[urr, "\imath_{\mathrm{p}}"] \arrow[r, "\operatorname{diag}_p"'] \arrow[drr, "\lambda_2"'] & X_p(G) \arrow[ur, "\pi_L"'] \arrow[dr, "\widetilde{\lambda}_{p,2}"] & \\
 & & \mathrm{PF}_2
\end{tikzcd}$$

(3)

Moreover, $\widetilde{\lambda}_{p,2}$ *is an algebra homomorphism.*

Proof. Since G is Kunze-Stein, it follows from (2) that there exists a bounded linear map $\imath_{\mathrm{KS}} : L^p(G) \to \mathrm{PF}_2(G)$, which makes the outer triangle in Diagram (3) commute. If $f \in \ker \imath_{\mathrm{KS}}$ then $f * h = 0$ for each $h \in C_c(G)$, so by basic measure theory $f = 0$ as an element of $L^p(G)$. Thus $\imath_{\mathrm{KS}}$ is injective.

Put $\widetilde{\lambda}_{p,2} := \imath_{\mathrm{KS}}\pi_L$; clearly this is continuous and linear, and it is injective since $\imath_{\mathrm{KS}}$ and π_L are. By construction , it makes the right-hand inner triangle in (3) commute. Now the top inner triangle in (3) commutes, by the definitions of diag_p and π_L. Hence, by a straightforward diagram chase, we find that the remaining inner triangle commutes.

Finally, since $\widetilde{\lambda}_{p,2} \operatorname{diag}_p = \lambda_2$ is a homomorphism and diag_p has dense range, it follows by continuity that $\widetilde{\lambda}_{p,2}$ is a homomorphism. ☐

Proof of Theorem 4.7. Let G be unimodular and Kunze-Stein. For $p = 2$, Theorem 4.7 is a special case of Theorem 3.6. So by Remark 4.1, it suffices to consider the case $1 \le p < 2$.

By Lemma 4.10, $\widetilde{\lambda}_{p,2} : X_p(G) \to \mathrm{PF}_2(G)$ is an *injective* algebra homomorphism; thus $X_p(G)$ is directly finite, using Theorem 3.6. Consider $\pi_P(X_p(G)) \subseteq \mathrm{PF}_p(G)$; this is directly finite, since π_P is an *injective* algebra homomorphism. On the other hand, by Lemma 4.9, $\pi_P(X_p(G))$ is a dense left ideal in $\mathrm{PF}_p(G)$. Therefore, by Proposition 2.7, we conclude that $\mathrm{PF}_p(G)$ is directly finite. ☐

5 TWO GROUPS FOR WHICH $L^1(G)$ IS NOT DIRECTLY FINITE

Let $\mathbb{F}$ be either $\mathbb{R}$ or $\mathbb{C}$, equipped with its usual topology: write $\mathbb{F}^\times$ for $\mathbb{F} \setminus \{0\}$, regarded as a multiplicative group. The affine group of $\mathbb{F}$, denoted by $\mathrm{Aff}(\mathbb{F})$, is defined to be the group

$$\left\{ \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} \;:\; a \in \mathbb{F}^{\times}, b \in \mathbb{F} \right\}$$

equipped with the natural topology.

The main result of this section is the following theorem.

THEOREM 5.1. *Neither* $L^1(\mathrm{Aff}(\mathbb{R}))$ *nor* $L^1(\mathrm{Aff}(\mathbb{C}))$ *are directly finite.*

REMARK 5.2. As remarked briefly in the introduction, λ_p extends to a continuous, *injective* algebra homomorphism $L^1(G) \to \mathrm{PF}_p(G)$ (continuity and injectivity follow from some standard measure-theoretic arguments). It therefore follows from Theorem 5.1 that if $G = \mathrm{Aff}(\mathbb{R})$ or $G = \mathrm{Aff}(\mathbb{C})$, none of the algebras $\mathrm{PF}_p(G)$ are directly finite.

Our approach to proving Theorem 5.1 is somewhat indirect. We first consider the group C*-algebras of $\mathrm{Aff}(\mathbb{R})$ and $\mathrm{Aff}(\mathbb{C})$, since we can use results of Diep and Rosenberg to show these algebras are not directly finite. Then, using the fact that these groups are Hermitian (see below for the definition), we deduce that the L^1-group algebras also fail to be directly finite. The ideas in this last step are taken from some work of Barnes [1], on spectra of convolution operators.

We start with some general definitions. A $*$-algebra A is said to be symmetric if $\sigma(x^*x) \subseteq [0,\infty)$ for all $x \in A$, and hermitian if $\sigma(h) \subseteq \mathbb{R}$ for all self-adjoint $h \in A$. (The spectrum is taken relative to A or to $A^\sharp$, depending on whether A has an identity element.) The two notions coincide for Banach $*$-algebras (see [18, Theorem 11.4.1]).

A locally compact group G is said to be hermitian if $L^1(G)$, equipped with the usual involution, is a hermitian Banach $*$-algebra. Further background on Hermitian groups can be found in [18, §12.6.22]; we only require the following examples, due to Leptin.

THEOREM 5.3 (Leptin). $\mathrm{Aff}(\mathbb{R})$ *and* $\mathrm{Aff}(\mathbb{C})$ *are Hermitian.*

Proof. This follows from a slightly more general result [13, Satz 6]. ☐

Barnes observed that groups which are both hermitian and amenable have the following spectral permanence property.

PROPOSITION 5.4 (Barnes). *Let G be a hermitian, amenable group, and let* $\theta : L^1(G) \to \mathcal{B}(\mathcal{H})$ *be a faithful* $*$*-representation on some Hilbert space. If* $h = h^* \in L^1(G)$*, then* $\sigma_{L^1(G)}(h) = \sigma_{\mathcal{B}(\mathcal{H})}\theta(h)$*.*

Proof. This is essentially contained in [1, Theorem 6]. ☐

REMARK 5.5. Strictly speaking, the statement of [1, Theorem 6] only gives $\sigma_{L^1(G)}(h) = \sigma_{C^*_r(G)}\lambda_2(h)$. However, examination of the proof, together with the theorem of Hulanicki that is quoted in [1, p. 329], shows that λ_2 can be replaced by any faithful $*$-representation of $L^1(G)$ on Hilbert space. The key point is that since G is amenable, the reduced and full group C*-algebras coincide.

CALCULATIONS OF DIEP AND OF ROSENBERG We first treat the case $G = \mathrm{Aff}(\mathbb{R})$, for which our main source is Chapter 3 of the monograph [6]. The results originally date from Diep's thesis, and are summarized without full proofs in [22].

The irreducible, continuous, unitary representations of $\mathrm{Aff}(\mathbb{R})$ were first worked out by Gelfand and Naimark. We focus on one in particular: let $\mathcal{H}_S$ be the Hilbert space $L^2(\mathbb{R}^\times, |x|^{-1}\,dx)$, and define a strongly continuous unitary representation $S : \mathrm{Aff}(\mathbb{R}) \to \mathcal{B}(\mathcal{H}_S)$ by

$$(S_g f)(x) = e^{ibx} f(ax) \qquad \text{for } f \in \mathcal{H}_S \text{ and } g = \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix}.$$

The representation S is quasi-equivalent to the left regular representation, and so in particular the induced $*$-homomorphism $\mathrm{C}^*(\mathrm{Aff}(\mathbb{R})) \to \mathcal{B}(\mathcal{H}_S)$ is injective. (See [6, proof of Lemma 3.3].)

The following result is proved in work of Diep, but may have been known earlier.

THEOREM 5.6 (Diep). *There exists* $f \in L^1(\mathrm{Aff}(\mathbb{R}))$, *such that* $I - S(f) : \mathcal{H}_S \to \mathcal{H}_S$ *is injective with closed range of codimension one.*

Proof. Define $h \in L^1(\mathrm{Aff}(\mathbb{R}))$ by

$$h\left(\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix}\right) = \chi_{|a|\leq 1} \frac{2a^2}{\sqrt{2\pi}} \exp(-b^2/2)\ . \tag{4}$$

By [6, Lemmas 3.6 and 3.7], $I - S(h)$ is invertible modulo the compacts, i.e. it is a Fredholm operator on $L^2(\mathcal{H}_S)$. In particular, it has closed range. Furthermore, by [6, Lemmas 3.9 and 3.10], $I - S(h)$ has 1-dimensional kernel and is surjective. We therefore take $f = h^* \in L^1(\mathrm{Aff}(\mathbb{R}))$; by the previous observations, $I - S(f) = (I - S(h))^*$ is injective with closed range, and has one-dimensional cokernel. ☐

REMARK 5.7. We have only taken from Diep's work what is needed for the present article. For a fuller discussion of how $\mathrm{C}^*(\mathrm{Aff}(\mathbb{R}))$ arises as an extension of an abelian C^*-algebra by $\mathcal{K}(\mathcal{H}_S)$, and the role played by the Fredholm operator $I - S(h)$ in classifying $\mathrm{C}^*(\mathrm{Aff}(\mathbb{R}))$, see [6, §3].

In the case $G = \mathrm{Aff}(\mathbb{C})$, we have a result analogous to Theorem 5.6, obtained by Rosenberg [20] using a suitable adaptation of Diep's methods. Let $\mathcal{H}_S$ be the Hilbert space $L^2(\mathbb{C}^\times, |z|^{-1}\,dz)$, and define a strongly continuous unitary representation $S : \mathrm{Aff}(\mathbb{C}) \to \mathcal{B}(\mathcal{H}_S)$ by

$$(S_g f)(z) = e^{i\,\mathrm{Re}\,wz} f(az) \qquad \text{for } f \in \mathcal{H} \text{ and } g = \begin{pmatrix} a & w \\ 0 & 1 \end{pmatrix}.$$

THEOREM 5.8 (Rosenberg). *There exists* $f \in L^1(\mathrm{Aff}(\mathbb{C}))$ *such that* $I - S(f) : \mathcal{H}_S \to \mathcal{H}_S$ *is injective with closed range of codimension one.*

We omit the example and proof, which can be found, modulo some small adjustments, in [20, Proposition 1].

Proof of Theorem 5.1. We first treat the case of $\mathrm{Aff}(\mathbb{R})$. Applying Theorem 5.6, there exists a faithful unitary representation $S : \mathrm{Aff}(\mathbb{R}) \to \mathcal{H}_S$, and some $f \in L^1(\mathrm{Aff}(\mathbb{R}))$, such that $I - S(f) : \mathcal{H}_S \to \mathcal{H}_S$ is injective with closed range of codimension one.

Since $I - S(f)$ is not invertible in $\mathcal{B}(\mathcal{H}_S)$, f is not quasi-invertible in $L^1(\mathrm{Aff}(\mathbb{R}))$. On the other hand, note that

$$I - S(f^* \bullet f) = I - S(f)^* \bullet S(f) = (I - S(f))^*(I - S(f)),$$

which *is* invertible in $\mathcal{B}(\mathcal{H}_S)$. Now, since $\mathrm{Aff}(\mathbb{R})$ is a Hermitian group, and S is faithful, by Proposition 5.4 we have

$$\sigma_{L^1(\mathrm{Aff}(\mathbb{R}))^\sharp}(f^* \bullet f) = \sigma_{\mathcal{B}(\mathcal{H}_S)}\phi(f^* \bullet f)\ .$$

In particular, $f^* \bullet f$ is quasi-invertible in $L^1(\mathrm{Aff}(\mathbb{R}))$, with quasi-inverse h, say. Then $h \bullet f^* \in L^1(\mathrm{Aff}(\mathbb{R}))$ and $h \bullet f^* \bullet f = 1$, showing that $L^1(\mathrm{Aff}(\mathbb{R}))$ is not directly finite.

The proof for $\mathrm{Aff}(\mathbb{C})$ is exactly similar, except that we use Theorem 5.8 instead of Theorem 5.6. ☐

6 CONCLUDING THOUGHTS

Theorems 3.6 and 5.1 immediately suggest the natural question:

QUESTION 1. *For which locally compact groups G is $\mathrm{C}^*_r(G)$ directly finite?*

Even in the special case where G is assumed to be a solvable Lie group, Example 3.7 suggests that a full characterization may be somewhat tricky to obtain.

For unimodular groups, we woud like to know if $\mathrm{PF}_p(G)$ is always directly finite. Note that the partial results in Section 4 do not apply if G is both discrete and non-amenable, and so we may in particular pose the following:

QUESTION 2. *Let $\mathbb{F}_2$ denote the free group on two generators. Does there exist $p \in (1, 2)$ for which $\mathrm{PF}_p(\mathbb{F}_2)$ is* not *directly finite?*

Since $\mathrm{PF}_p(\mathbb{F}_2)$ contains as a dense subalgebra the directly finite algebra $\ell^1(\mathbb{F}_2)$, a positive answer to Question 2 would give a "natural", negative solution to our final question.

QUESTION 3. *Is the completion of a directly finite normed algebra itself directly finite?*

We suspect not, but know of no counterexample. On the other hand, if Question 3 has a *positive* answer, then $\mathrm{PF}_p(G)$ is directly finite for every unimodular group and all $p \in (1, \infty)$.

ACKNOWLEDGEMENTS

The present paper has a somewhat tangled history. The seeds for it (in particular, the observations that became Section 2, Proposition 3.1 and Corollary 4.4) were sown during the author's time as a postdoctoral researcher in the Département de mathématiques et de statistique at Université Laval, whose support is gratefully acknowledged. Further developments, in particular the discovery via [20] that certain reduced group C*-algebras are not directly finite, were made during a visit to the University of Leeds in May 2010, as part of a working semester on *Banach algebra and operator space techniques in topological group theory.* The author thanks the University of Leeds for its hospitality.

Other main results (such as the Kunze-Stein case, and the proof that certain L^1-group algebras are not directly finite), location of various important references, and the final write-up, all date from the author's current position at the University of Saskatchewan. This activity was supported by a start-up grant from the University, and also by NSERC Discovery Grant 402153-2011.

Thanks are due to various colleagues and correspondents, in particular: J. Meyer, for pointing out the reference [11]; E. Samei, for providing the reference [5]; and M. Daws, for useful conversations regarding the basic theory of Hilbert algebras.

A INGREDIENTS IN THE PROOF OF THEOREM 4.3

To fix notation and provide background, we quickly summarize the necessary definitions from [9]. Throughout, $p \in (1,\infty)$ and q is the conjugate index to p.

Fix a left Haar measure μ on G. Define a contractive linear map $\theta_p : L^p(G) \mathbin{\widehat{\otimes}} L^q(G) \to C_0(G)$ by

$$\theta_p(f\otimes g)(x) = \langle \lambda_p(x)f, g\rangle = \int_G f(x^{-1}y)g(y)\,d\mu(y) \qquad (f\in L^p(G), q\in L^q(G), x\in G),$$

and let $\mathrm{A}_p(G) \subseteq C_0(G)$ be the *coimage* of θ_p (more explicitly, the image of θ_p equipped with the quotient norm induced from $\left(L^p(G) \mathbin{\widehat{\otimes}} L^q(G)\right) / \ker\theta_p$). (It can be shown, although we do not need this, that the spaces $\mathrm{A}_p(G)$ are subalgebras of $C_0(G)$, called the Figa-Talamanca–Herz algebras.) If we restrict θ_p to the dense subspace $C_c(G)\otimes C_c(G)$, we obtain a linear map $\theta : C_c(G)\otimes C_c(G) \to C_c(G)$, whose image we denote by $\mathcal{A}_c(G)$. Note that $\mathcal{A}_c(G)$ is dense in $\mathrm{A}_p(G)$, for every p.

Integration on G defines a pairing $C_c(G)\times C_0(G) \to \mathbb{C}$ and hence we have natural maps $C_c(G) \to C_0(G)^* \to \mathrm{A}_p(G)^*$, the second map being restriction. We may thus regard θ_p^* as a map $C_c(G) \to (L^p(G) \mathbin{\widehat{\otimes}} L^q(G))^* \cong \mathcal{B}(L^p(G))$, which can be described explicitly as follows: given $f\in C_c(G)$ and $\xi\in L^p(G)$, $\eta \in L^q(G)$, we have

$$\langle \theta_p^*(f)\xi, \eta\rangle = \int_G f(t)\theta_p(\xi\otimes\eta)(t)\,d\mu(t)$$

Moreover, if $\xi,\eta\in C_c(G)$ then it is easily shown by rearranging the integral that $\langle\theta_p^*(f)\xi,\eta\rangle = \langle\lambda_p(f)\xi,\eta\rangle$, and since both $\theta_p^*(f)$ and $\lambda_p(f)$ are bounded operators we have$\theta_p^*(f) = \lambda_p(f)$ by density. In particular, $\lambda_p(f) \in \theta_p^*(\mathrm{A}_p(G))$ for all $f\in C_c(G)$.

Note that since $\theta_p^* : \mathrm{A}_p(G)^* \to \mathcal{B}(L^p(G))$ is the adjoint of a quotient map, it is an isometry with weak-star closed range. It follows that $\theta_p(\mathrm{A}_p(G))^* \supseteq \mathrm{PF}_p(G)$, and so there is an isometry $\phi_p : \mathrm{PF}_p(G) \to \mathrm{A}_p(G)^*$, such that $\theta_p^*\phi_p$ is just the inclusion of $\mathrm{PF}_p(G)$ into $\mathcal{B}(L^p(G))$.

The key result needed from [9] is the following theorem, whose proof we omit. It should be emphasised that the theorem is independent of the results in [10].

THEOREM A.1 ([9, Theorem 1]). *Let G be a locally compact group and let $p \in (1,\infty)$. Then the map $\mathrm{A}_p(G) \mathbin{\widehat{\otimes}} \mathrm{A}_2(G) \to C_0(G)$ defined by multiplication of functions takes values in $\mathrm{A}_p(G)$, and*

$$\|fh\|_{\mathrm{A}_p(G)} \leq \|f\|_{\mathrm{A}_2(G)} \|h\|_{\mathrm{A}_p(G)} \tag{5}$$

for all $f \in \mathrm{A}_p(G)$ and $h \in \mathrm{A}_2(G)$.

We also need a result that is stated and proved in [10, §9], where it is described as being folklore.

LEMMA A.2. *Let G be amenable and let $p \in (1,\infty)$. Given any compact set $K \subseteq G$, and any $\varepsilon > 0$, there exists a compactly supported function $f \in \mathrm{A}_p(G)$ which is identically 1 on K and has $\mathrm{A}_p(G)$-norm at most $1+\varepsilon$.*

Proof of Theorem 4.3. Let $h \in \mathrm{A}_2(G)$ have compact support. Taking $K = \operatorname{supp}(h)$ in Lemma A.2 and using (5), we get $h \in \mathrm{A}_p(G)$ and $\|h\|_{\mathrm{A}_p(G)} \leq (1+\varepsilon)\|h\|_{\mathrm{A}_2(G)}$, for all $\varepsilon > 0$. Since $\mathrm{A}_c(G)$ is norm-dense in $\mathrm{A}_2(G)$, this implies that $\mathrm{A}_2(G) \subseteq \mathrm{A}_p(G)$ with non-increase of norms. Moreover, since $\imath(\mathcal{A}_c(G)) = \mathcal{A}_c(G)$, and $\mathcal{A}_c(G)$ is dense in $\mathrm{A}_p(G)$, the inclusion $\imath : \mathrm{A}_2(G) \to \mathrm{A}_p(G)$ has dense range. Hence $\imath^* : \mathrm{A}_p(G)^* \to \mathrm{A}_2(G)^*$ is injective, and has norm ≤ 1.

Now consider the following commutative diagram:

$$\begin{tikzcd}
C_c(G) \arrow[r, "\lambda_p"] \arrow[dr, "\lambda_2"'] & \mathrm{PF}_p(G) \arrow[r, "\phi_p"] & \mathrm{A}_p(G)^* \arrow[d, "\imath^*"] \\
& \mathrm{PF}_2(G) \arrow[r, "\phi_2"'] & \mathrm{A}_2(G)^*
\end{tikzcd}$$

We have, for each $f \in C_c(G)$,

$$\|\lambda_2(f)\| = \|\phi_2\lambda_2(f)\| = \|\imath^*\phi_p\lambda_p(f)\| \leq \|\lambda_p(f)\|.$$

Therefore, since $\lambda_p(C_c(G))$ is norm-dense in $\mathrm{PF}_p(G)$, there is a unique continuous linear map $J : \mathrm{PF}_p(G) \to \mathrm{PF}_2(G)$ such that $J\lambda_p = \lambda_2$. By continuity, J is an algebra homomorphism. Moreover, for each $f \in C_c(G)$,

$$\phi_2 J\lambda_p(f) = \phi_2\lambda_2(f) = \imath^*\phi_p\lambda_p(f)$$

so by density and continuity, $\phi_2 J = \imath^*\phi_p$. Since $\imath^*\phi_p$ is injective, so is J, and the proof of Theorem 4.3 is complete. □

REFERENCES

[1] B. A. BARNES, *When is the spectrum of a convolution operator on L^p independent of p?*, Proc. Edinburgh Math. Soc. (2), 33 (1990), pp. 327–332.

[2] F. F. BONSALL AND J. DUNCAN, *Complete normed algebras*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 80, Springer-Verlag, New York, 1973.

[3] Y. CHOI, *Group representations with empty residual spectrum*, Int. Eq. Op. Th., 67 (2010), pp. 95–107. Erratum: Int. Eq. Op. Th. 69 (2011), no. 1, 149–150.

[4] M. COWLING, *The Kunze-Stein phenomenon*, Ann. Math. (2), 107 (1978), pp. 209–234.

[5] A. DERIGHETTI, *Convolution operators on groups*, vol. 11 of Lecture Notes of the Unione Matematica Italiana, Springer, Heidelberg, 2011.

[6] D. N. DIEP, *Methods of noncommutative geometry for group C^*-algebras*, vol. 416 of Chapman & Hall/CRC Research Notes in Mathematics, Chapman & Hall/CRC, Boca Raton, FL, 2000.

[7] J. DIXMIER, *Les algèbres d'opérateurs dans l'espace hilbertien (algèbres de von Neumann)*, Gauthier-Villars Éditeur, Paris, 1969. Deuxième édition, revue et augmentée, Cahiers Scientifiques, Fasc. XXV.

[8] ——, *C^*-algebras*, North-Holland Publishing Co., Amsterdam, 1977. Translated from the French by Francis Jellett, North-Holland Mathematical Library, Vol. 15.

[9] C. HERZ, *The theory of p-spaces with an application to convolution operators.*, Trans. Amer. Math. Soc., 154 (1971), pp. 69–82.

[10] ——, *Harmonic synthesis for subgroups*, Ann. Inst. Fourier (Grenoble), 23 (1973), pp. 91–123.

[11] I. KAPLANSKY, *Modules over operator algebras*, Amer. J. Math., 75 (1953), pp. 839–858.

[12] R. A. KUNZE AND E. M. STEIN, *Uniformly bounded representations and harmonic analysis of the* 2×2 *real unimodular group*, Amer. J. Math., 82 (1960), pp. 1–62.

[13] H. LEPTIN, *Lokal kompakte Gruppen mit symmetrischen Algebren*, in Symposia Mathematica, Vol. XXII (Convegno sull'Analisi Armonica e Spazi di Funzioni su Gruppi Localmente Compatti, INDAM, Rome, 1976), Academic Press, London, 1977, pp. 267–281.

[14] J. MEYER, *personal communication*. MathOverflow. `http://mathoverflow.net/questions/16944` (version: 2010-03-09).

[15] M. S. MONTGOMERY, *Left and right inverses in group algebras*, Bull. Amer. Math. Soc., 75 (1969), pp. 539–540.

[16] W. D. MUNN, *Direct finiteness of certain monoid algebras. I*, Proc. Edinburgh Math. Soc. (2), 39 (1996), pp. 365–369.

[17] C. NEBBIA, *Groups of isometries of a tree and the Kunze-Stein phenomenon*, Pacific J. Math., 133 (1988), pp. 141–149.

[18] T. W. Palmer, *Banach algebras and the general theory of $*$-algebras. Vol. 2*, vol. 79 of Encyclopedia of Mathematics and its Applications, Cambridge University Press, Cambridge, 2001.

[19] G. K. Pedersen, *C^*-algebras and their automorphism groups*, vol. 14 of London Mathematical Society Monographs, Academic Press Inc. [Harcourt Brace Jovanovich Publishers], London, 1979.

[20] J. Rosenberg, *The C^*-algebras of some real and p-adic solvable groups*, Pacific J. Math., 65 (1976), pp. 175–192.

[21] X. Wang, *The C^*-algebras of a class of solvable Lie groups*, vol. 199 of Pitman Research Notes in Mathematics Series, Longman Scientific & Technical, Harlow, 1989.

[22] D. N. Z′ep, *The structure of the group C*-algebra of the group of affine transformations of the line*, Funkcional. Anal. i Priložen., 9 (1974), pp. 63–64.

Department of Mathematics and Statistics
McLean Hall
University of Saskatchewan
106 Wiggins Road, Saskatoon, SK
Canada, S7N 5E6
Email: `choi@math.usask.ca`